

EMPTY NOSE SYNDROME

A Guide to Diagnosis and Management
for Medical Professionals



EMPTY NOSE SYNDROME
INTERNATIONAL
ASSOCIATION

BACKGROUND

As a surgical complication secondary of turbinate surgery, Empty Nose Syndrome is an uncommon condition, which is underdiagnosed and is frequently untreated. There is a common misperception among some medical professionals that ENS is not severe or substantially disabling and can be managed with conservative treatments. This perception has resulted in ENS being little studied and largely overlooked, with a general lack of proper attention in both research into its cause and, more importantly, into its cure.

The purpose of this document is to provide guidance for medical professionals concerning diagnosis, management, and treatment of ENS.

The information contained herein is based on current medical literature and experience with ENS patients. The information should be considered advisory only.

ABOUT ENS

History

The term Empty Nose Syndrome was first coined by Dr. Eugene Kern of the Mayo Clinic in 1994. Dr. Kern described the condition and studied patients, many of whom had experienced total turbinectomies, a procedure which had seen a resurgence in the 1970's (lasting through the 1990s) having been earlier abandoned following condemnation around the turn of the century. The condition in connection with turbinate procedures, especially the turbinectomy, was known by the late 19th century (described as atrophy).

Definition

Empty Nose Syndrome is an iatrogenic disorder - a complication of nasal surgery - where the nasal turbinates, especially the inferior turbinates, have been excised or damaged as a result of turbinate surgery causing the destruction of normal nasal physiology. This leads to clinical symptoms of paradoxical obstruction *, nasal dryness, crusting, intermittent bleeding, lack of airflow sensation, nasal and upper respiratory dryness/irritation/pain from unprocessed/cold/dry inspired air, disruption in nervous system function, and number of secondary symptoms. In brief, ENS can be considered a nasal breathing

* Feeling of nasal obstruction is described as constant and continuous: feeling of suffocation, inability or significant difficulty to breathe through nose, feeling that nose is too open, sensation of excessive airflow, shortness of breath, difficulty to properly inflate the lungs, lack of nasal resistance, and/or undifferentiated breathing difficulties.

disorder (characterized by paradoxical obstruction) along with concomitant nasal dryness (atrophy) ^{1 2}.

Statistics and Incidence rate

Empty Nose Syndrome affects a small number of the population and the incidence is not known because there are no specific studies which have investigated the incidence of ENS. The absence of incidence-related studies is directly related to the lack of awareness about ENS among medical professionals. This has resulted in the absence of diagnostic criteria and omission of a diagnosis of ENS in patient records.

Causes

ENS occurs following turbinate surgery and can emerge months to years afterward. Any turbinate procedure has the potential to cause ENS if it is performed too aggressively. Some turbinate procedures increase the likelihood of ENS, such as partial or total resection of the inferior nasal turbinates, or mucosal surface cautery ³. Risk of ENS increases the greater the turbinates are excised or damaged. Total turbinectomies are less common today following the resurgence of the 1970's and are frowned upon, especially within the specialty of otorhinolaryngology, however, there is no general prohibition against them.

Currently, there is no consensus concerning how much turbinate excision or damage can result in ENS, nor what measure of excision or damage constitutes an “aggressive” reduction. On the subject of ‘how much is too much?’, there exist no quantifying studies, guidelines, or formal statements issued by professional bodies.

Features of ENS

ENS shares features in common with secondary atrophic rhinitis, but current research has distinguished it as a separate condition. Despite ENS lacking a diagnostic consensus, most ENTs consider ENS to be a true clinical entity, and its primary features are agreed upon. ENS is a multifactorial condition, which arises due to a combination of structural and functional impairments of the vital nasal structures, involving altered nasal aerodynamics, reduced and impaired mucosal area, and neurosensory damage (e.g. damage to cool thermoreceptors) ⁴.

ENS patients have significant difficulties including:

- constant dyspnea and/or a sensation of suffocation;
- sleep deprivation;
- nasal, facial, and respiratory dryness, coldness, irritation, and pain;
- increased illnesses from damaged nasal physiology such as recurrent sinus infection, bronchitis, asthma;
- a disrupted nasal cycle;
- autonomic nervous system dysfunction;
- a variety of other medical complications secondary to ENS.

ENS is a debilitating and disabling condition, resulting in a poor quality of life since good nasal function is crucial for proper lung functions and breathing, body’s oxygenation, ability to function physically, proper functioning of nervous system, cognitive functions, and sense of well-being. ENS patients suffer substantially and continuously.

Mouth breathing is not a reasonable alternative, since it is not a physiological method of respiration and should never be considered a substitute for nasal breathing. Nasal airway resistance accounts for more than 50% of total airway resistance and increases oxygen (O₂) uptake by 10-20% ⁵. Mouth breathing does not alleviate ENS symptoms.

Diagnosis

Formerly, the difficulty has been in obtaining objective diagnostic evidence, rather than subjective symptomatology of patients (derived, for example, from the Sino Nasal Outcome Test, SNOT). Even so, conditions such as tinnitus are diagnosed despite there being a similar impediment to diagnosis. However, there are a number of diagnostic tools, some of which are new, some of which have not been routinely employed in the diagnosis of ENS. ENS is currently able to be diagnosed within the clinical setting, and diagnosis is able to be further supported by diagnostic tools available outside the clinical setting.

Treatment, Prognosis, Quality of Life

Conservative management of ENS focuses on irrigation and moisturizing of the nose to preserve remaining mucosa and needs to be performed for the patient’s lifetime. The necessity for chronic nasal care is burdensome and time consuming.

While nasal reconstructive surgery and regenerative medicine treatments can result in some improvement in symptoms, it is important to remember that turbinate tissue cannot be replaced or regrown, and there is no cure for ENS.

ENS patients experience a significant reduction in quality of life (QOL). ENS has an impact on physical health, employment, marital, social, and financial aspects of the patient's life. As a chronic illness, ENS can have an impact on the patient's well-being and reduce his/her enjoyment of life, inducing stress and secondary symptoms like anxiety and depression. ENS can cause inability to work or significant number of 'sick days missed from work, decreased productivity, and lifestyle disruption⁶. There are memory and concentration deficits caused by abnormal breathing, the lack of sleep, a diminished sense of smell, and lack of nasal sensation which also lower QOL.

ENS produces environmental limitations for the patient in windy, dry, cold, hot, air-conditioned, heated, dusty or odor-filled places, because the nose can no longer properly fulfill its functions (filtering, humidifying, dehumidifying, warming, smelling, and safeguarding).

To place the symptoms and sequelae of ENS in context, a diminished sense of smell (hyposmia) in and of itself can be considered a major contributor to reduced QOL.

These difficulties in their total can result in the ENS patient suffering from general disability, that is, a difficulty functioning and performing daily activities.

To assess of quality of life of ENS patient it is advisable to use the **Rhinosinusitis Disability Index (RSDI)**. Another generic assessment instrument for health and disability is the **WHO Disability Assessment Schedule**⁷.

Statistics

ENS is considered a rare condition, affecting a small subset of patients who undergo turbinate reduction. However, the true incidence is unknown since there have been no specific studies published investigating the incidence. This is due to the fact that there is a lack of information concerning the condition and no universal definitional/diagnostic consensus. A 22.2% incidence of atrophy following inferior turbinectomy was reported in a long-term study⁸. Other estimates say that 20% of patients with resection will develop fulminant ENS and a greater percentage will suffer dryness or changes in airflow⁹.

The most recent data from 2006 suggest there are an estimated 196,000 turbinectomies in the United States according to National Center for Health Statistics¹⁰. Turbinectomies often are performed as secondary procedures in conjunction with surgeries such as functional endoscopic sinus surgery (FESS) and/or a septoplasty. Since this figure does not include turbinectomies performed in conjunction with other sino-nasal surgeries, the number of turbinate surgeries is likely much higher¹¹.

In addition, a portion of the middle turbinates may be resected during FESS or other procedures in order to facilitate access to certain areas of the nose.

Clinical presentation

Nasal dryness and breathing difficulties are the most common presenting symptoms with paradoxical obstruction being considered the hallmark symptom. Patients complain of stuffiness or emptiness despite having a patent nasal cavity. They may also describe a feeling of lack of nasal resistance, difficulty to properly inflate the lungs, breathlessness, suffocation, or lack of nasal airflow sensation. Nasal dryness presents on a continuum ranging from simple dryness to atrophy. Mucosa can appear pale and dry. There may be thick or absent secretions. Thick secretions can cause congestion, pressure,

and pain. Crusting is often present, with significant crusting in more severe cases. Additionally, there may be significant nasal and facial pain caused by mucosal dryness, atrophy, or neural damage. Hyposmia and hypersensitivity to volatile compounds ** can both affect the patient's sense of smell and quality of life. Symptoms can range from mild to extremely severe.

Patients presenting with ENS can have a substantial loss of mucosal turbinate tissue, or can have normal appearing turbinates and intranasal volume (having suffered functional damage to the turbinates). Clinical presentation within the ENS population varies. Patients upon nasal examination may vary in appearance due to the method and amount of turbinate reduction. This is due to the fact that it is not known how much turbinate tissue can be removed or damaged before nasal physiology is compromised or before the aerodynamic pattern of the nasal cavity becomes impaired. Symptomatology may vary in number of symptoms and severity, but the well-established primary symptoms of ENS are often universally present, the core of which comprise Houser's five modifications to the SNOT-25. Since the onset of ENS can occur at an interval ranging from weeks, months, to years, the patient's turbinate procedure may have been recent or taken place some years prior.

** Volatile compounds - gasoline, perfume, smoke, smells, etc.

ENS Subtypes

ENS has been classified into four subtypes: ENS-IT, ENS-MT, ENS-Both, and ENS-Type¹².

The subtypes describe the following:

- IT - where the inferior turbinate has been partially or totally excised;
- MT - where the middle turbinate has been partially or fully excised;
- Both - where the inferior and middle turbinates have been at least partially excised or reduced;
- Type - where a patient has sufficient-appearing turbinate tissue, but suffers ENS symptoms after surgery affecting the mucosal surface of the turbinates¹³.

Complicating Factors

Common findings of complicating factors may include:

- additional nasal openness from concomitant septoplasty (which can affect/diminish the septal swell body);
- septal perforations;
- structural collapse, such as nasal valve collapse;
- large sinus openings and absent nasal structures from overly aggressive sinus or nasal surgery that facilitate unphysiological ventilation;
- mucosa affected by age, autoimmune disease or hormonal status (e.g. geriatric rhinitis, Sjogren's syndrome, and menopause).

It is important to inquire of the patient about his moisturizing habits. Use of humidifiers, saline, and emollients may interfere with examination-based evaluation of nasal dryness. A nose with healthy physiology does not require any moisturizing.

A Note on Patients Presenting with ENS

Patients presenting with ENS for clinical examination are often in distress. They will commonly have been suffering for a considerable length of time without relief from severe breathing difficulties, damaged sensory ability, severe nasal dryness with its attendant acute burning and pain, and from acute sleep deprivation. The majority of patients have no understanding of nasal physiology and are unfamiliar with related symptoms. They often have difficulty expressing symptoms. Among the most difficult symptoms is *paradoxical obstruction*. Patients with significantly resected turbinates may suffer from lack of nasal resistance necessary for efficient respiration and pulmonary gas exchange. Additionally, patients may experience lack of airflow sensation, which creates conflict in signals between their nose and central nervous system (brain). This results in a sense of suffocation. Patients feel that they have lost control of their breathing, sleep, and important sensory input (airflow, temperature, smell), in addition to the pain of severe nasal dryness and unprocessed airflow in the upper respiratory tract due to excessive nasal openness. This situation often results in a heightened state of anxiety and may trigger a continuous “fight-or-flight” response in patients.

Table 1: lists common patient complaints:

- Nose feels too open/empty
- Getting too much air (excessive airflow)
- Breathlessness and/or feeling of suffocation
- Feeling of hyperventilation
- Difficulty to inflate lungs/draw a full breath
- Lack of airflow sensation
- Cold, dry, unprocessed air hitting the back of throat
- Severe dryness with/without crusts
- Burning/pain in nose
- Nose feels numb
- Unable to sleep because of difficulty breathing
- Cannot find right way to breathe through the nose
- Unable to relax/feel like nervous system is going “crazy”

SYMPTOMS

ENS symptomatology has been assessed utilizing the SNOT 20 or 22 (Sino-nasal Outcome Test). Subsequently, the SNOT-25 has been utilized to assess ENS symptomatology¹⁴. The new ENS6Q (Empty Nose Syndrome 6 Questionnaire) has also been used for assessment.

ENS-specific questions are contained in both the SNOT-25 and the ENS6Q. The SNOT-25 contains five (5) items specific to ENS¹⁵. The ENS6Q is a validated ENS-specific, 6-item questionnaire, which can be used as an adjunct to the SNOT-22 to screen patient for ENS¹⁶. The ENS6Q includes four (4) of the ENS-specific items from the SNOT-25, adding two new items: sense of diminished airflow (cannot feel air flowing through your nose) and nasal burning.

SNOT-55

ENSIA has proposed a fifty-five (55) item modified SNOT test to assess ENS symptoms. This proposed SNOT-55 builds on SNOT-25 adding thirty (30) new items.

These new items define symptoms that have been widely discussed and thoroughly described by ENS sufferers in support groups over the past 10 years.

Some of these new items have previously been described in the works of SM Houser, EB Kern, and in the book of patient (and ENS sufferer) Christopher Martin, but have heretofore not been applied specifically in a test or measure¹⁷.

The thirty questions are scored from 0-5 with higher scores indicating greater ENS-related disease burden (health concerns) including physical & emotional problems and functional limitations.

Description of items

Items 1-25 comprise the SNOT-25. A large number of new items in the SNOT-55 are related to the 'breathing disorder' component of ENS. These include items 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 40, & 54. Breathing-related symptoms vary among ENS patients and can differ based on the method and extent of turbinate reduction.

The number of breathing-related items contained in SNOT-55 reflect this variation. For example, a patient with a large resection may experience breathing difficulties differently than someone who has seemingly adequate turbinate tissue that has experienced surface cautery. Additional rhinological symptoms are included as items 41 through 47. Other secondary physical symptoms are listed as items 48 through 50. Cognitive and functional limitations are expanded upon in items 51 through 55.

Table 2: SNOT-55

(ENS-specific symptoms from #21-55)

No.	SNOT – 55
Nasal Symptoms:	
1	Need to blow nose
2	Sneezing
3	Runny nose
4	Cough
5	Postnasal discharge
6	Thick nasal discharge (<i>thick mucus</i>)
7	Ear fullness
8	Dizziness
9	Ear pain
10	Facial pain/pressure
11	Difficulty falling asleep
12	Waking up at night
13	Lack of good night's sleep
14	Waking up tired
15	Fatigue
16	Reduced productivity
17	Reduced concentration
18	Frustration/restlessness/irritability
19	Sadness
20	Embarrassment
Houser Modification adds:	
21	[Nasal] dryness
22	Difficulty with nasal breathing
23	Suffocation
24	Nose is too open
25	Nasal crusting

ENSIA Modification adds:	
26	Nasal obstruction (<i>blockage /congestion of nose</i>)
27	Nasal emptiness
28	Excessive airflow (' <i>too much air</i> ')
29	Decreased nasal resistance
30	Shortness of breath (<i>dyspnea</i>)
31	Difficulty to inflate lungs
32	Difficulty drawing a full breath
33	Disturbed/disorganized airflow
34	Decreased sense of nasal airflow
35	Weakened airflow (<i>weak stream of inhaled air</i>)
36	Tight, asthma-like breathing
37	Use of accessory muscles (neck and chest) to facilitate breathing
38	Respiratory dryness/coldness/irritation of pharynx & adjacent areas (<i>breathing unprocessed air</i>)
39	Hypersensitivity to volatile compounds (e.g. <i>gasoline, perfume, smoke, smells, etc.</i>)
40	Rapid breathing
41	Reduced sense of smell
42	Lack of mucus
43	Nasal burning/pain
44	Nasal bleeding
45	Nasal inflammation
46	Impaired nasal drainage
47	Difficulty removing thick mucus/clearing nose
48	Dry eyes
49	Headache
50	Increased heart rate
51	Impaired concentration from breathing difficulties (<i>aproxexia nasalis</i>)
52	Anxiety
53	Increase in symptoms in windy, dry, cold, hot, air-conditioned, heated environments
54	Persistence of symptoms when mouth breathing
55	General disability (<i>difficulty functioning, performing daily activities</i>)

Table 3: ENS Symptoms with Definitions

Each symptom is provided with a definition.

<p>21 [Nasal] Dryness</p> <p>The consequence of the decompensation of the humidifying function of the nasal mucosa due to absent or dysfunctional turbinate tissue. Nasal mucosa can appear dry upon exam, or even atrophic.</p>	<p>22 Difficulty with nasal breathing</p> <p>Nasal breathing is made difficult by the changes in the nasal cavity following turbinate resection including: neurosensory damage, impaired aerodynamics, decreased nasal resistance, paradoxical obstruction from an overly patent airway. Patients can suffer from some or all of these changes.</p>	<p>23 Suffocation</p> <p>A sensation of suffocation is produced by a lack of nasal resistance necessary for efficient respiration & pulmonary gas exchange, and a lack of airflow sensation, which creates conflict in signals between the nose and central nervous system (brain).</p>	<p>24 Nose is too open</p> <p>Refers to a sensation that the patient's nose is too open due to lack of turbinate tissue. Patients may feel their nose is too open to airflow or to the environment. There can a sensation of excessive openness even while at rest which conveys a sense of vulnerability or lack of protection to the patient. Anxiety is a common response to this symptom.</p>
<p>25 Nasal crusting</p> <p>Hardened crusts that can develop in the dry nose from dried mucus, sores, and chronic bacteria. They can be painful and affect breathing.</p>	<p>26 Nasal obstruction <i>(blockage / congestion of nose)</i></p> <p>The feeling the nose is obstructed. In context of ENS, it is paradoxical obstruction – sensation of nasal obstruction despite a widely patent airway.</p>	<p>27 Nasal emptiness</p> <p>The sensation of absence or emptiness of the nose. Refers to the lack of nasal sensation due to removal of nasal structures and/or nasal neurosensory damage.</p>	<p>28 Excessive airflow (“too much air”)</p> <p>Refers to a sensation that the patient is breathing too much air, or breathing too freely.</p>
<p>29 Decreased nasal resistance</p> <p>Decreased values of nasal resistance in both inspiration and expiration occur in ENS. Adequate inspiratory resistance is necessary for efficient respiration and efficient gas-exchange in the alveoli of the lungs. Experienced in ENS as an absence of necessary “force” for adequate respiration.</p>	<p>30 Shortness of breath (dyspnea)</p> <p>Dyspnea. A feeling that one cannot breathe adequately. Can be related to a feeling of suffocation.</p>	<p>31 Difficulty to inflate lungs</p> <p>The feeling there is difficulty to inflate the lungs upon nasal inspiration. It is a feeling of shallow breathing. Due to changes in the nasal cavity - nasal resistance, airflow impairment.</p>	<p>32 Difficulty drawing a full breath</p> <p>Refers to the feeling of inspiration being limited to the upper chest, lacking diaphragmatic involvement. Due to decreased nasal resistance affecting pulmonary function. Impaired lung compliance may also be involved – such as bronchospasm due to inspiration of unconditioned air.</p>
<p>33 Disturbed/disorganized airflow</p> <p>A feeling of disorganized, disturbed airflow when breathing. Caused by changes in the nasal cavity following surgery.</p>	<p>34 Decreased sense of nasal airflow</p> <p>A phenomenon where the sensation of breathing is decreased or absent. Caused by damaged or lack of sensing nerve receptors of the inferior turbinate. Airflow can be sensed in the lungs, but not in the nose</p>	<p>35 Weakened airflow <i>(weak stream of inhaled air)</i></p> <p>Refers to the sensation that there is a weakened stream of airflow at a decreased velocity with inspiration. The sensation extends to a perception of shallow breathing.</p>	<p>36 Tight, asthma-like breathing</p> <p>Asthma-like symptoms such as bronchospasm and airway stenosis occurring as a result of inspiring unconditioned/unprocessed air and/ or lack of nasal resistance.</p>

<p>37 Use of accessory muscles (neck & chest) to facilitate breathing Use accessory muscles of the neck and chest to produce force and resistance within the upper airway & thoracic areas in order to facilitate breathing.</p>	<p>38 Respiratory dryness/coldness/irritation of pharynx & adjacent areas (<i>breathing unprocessed air</i>) A sensation of dryness, coldness, irritation in the pharynx and adjacent areas, especially noticeable when inspiring air. From the inspiring of unconditioned, unprocessed air. A central patient complaint indicative of ENS is dry, cold air hitting the back of the throat when breathing.</p>	<p>39 Hypersensitivity to volatile compounds (<i>e.g. gasoline, perfume, smoke, smells, etc.</i>) Refers to sensitivity to volatile compounds despite having reduced sense of smell generally. Possible reasons are changes in airflow, nasal emptiness, and dryness. The patient may experience nausea and irritant reactions.</p>	<p>40 Rapid breathing Rapid breathing can be either shallow or deep. Decreased nasal resistance can result in shallow breathing and increased breaths per minute. Decreased nasal airflow sensation may produce deep breathing as the patient struggles to sense adequate air intake, placing themselves in a state of hyperventilation. Both can occur under different circumstances and is also influenced by the type of turbinate reduction and surgical alterations of the nasal cavity as well as patient compensatory response to breathing impairment. Breathing at rest typically results in shallow breathing, while breathing with exertion or under stress produces the latter.</p>
<p>41 Reduced sense of smell Hyposmia - a reduced ability to smell and to detect odors. May be caused by altered airflow patterns, in particular lack of high inspiratory airflow, affecting olfactory stimulation/perception. Some ENS patients can suffer from anosmia.</p>	<p>42 Lack of mucus Turbinate tissue is rich in nasal glands. Absent, not sufficiently stimulated (due to decreased pressure and shear stress conditions as a result of resection of nasal structures), and/or damaged turbinate tissue results in a lack of mucus and nasal dryness. The important function of mucociliary clearance (MCC) is damaged.</p>	<p>43 Nasal burning/pain Burning sensation and pain within the nasal cavity. Can be both at rest and with breathing exacerbated by airflow. Dry nasal mucosa, inflammation, and nerve sensitivity contribute to this symptom.</p>	<p>44 Nasal bleeding Epistaxis. Caused by dry, fragile, or atrophied nasal mucosa.</p>
<p>45 Nasal inflammation Nasal inflammation caused nasal dryness, decreased mucus, impaired or resected nasal glands and vasculature of the turbinates.</p>	<p>46 Impaired nasal drainage Nasal drainage is impaired due to structural changes within the nasal cavity. The mucociliary clearance (MCC) system is damaged.</p>	<p>47 Difficulty removing thick mucus/clearing nose Impaired nasal aerodynamics, unusually thick mucus, impaired mucociliary clearance, nasal dryness make blowing or clearing the nose difficult.</p>	<p>48 Dry eyes Dry eyes can occur as a result of chronic nasal dryness.</p>

<p>49 Headache Breathing difficulties can contribute to headaches; more so due to hyperventilation¹⁸ or hypoxia. ENS-related sino-nasal symptoms contribute as well.</p>	<p>50 Increased heart rate Heart rate increased from the patient's normal rate. Need not reach tachycardia according to the "conventional rate limits for sinus rhythm"¹⁹, but may be a pathology for the individual patient^{***}. Caused by breathing difficulties, dyspnea or feeling of suffocation, and lack of nasal sensation (triggering autonomic stress from lack of signaling to the breathing centers of the brain).</p>	<p>51 Impaired concentration caused by breathing difficulties (<i>aproxexia nasalis</i>) More broadly can mean impaired cognitive function. Two descriptions of <i>aproxexia</i> from 19th century medical literature noted: "functional disorder of the brain, caused by disturbed nasal respiration" and "want of concentration".</p>	<p>52 Anxiety Physically generated anxiety is that induced by breathing difficulties, decreased airflow sensation, and autonomic disturbances.</p>
<p>53 Increase in symptoms in windy, dry, cold, hot, air-conditioned, heated environments Patients experience an increase in symptoms both indoors and outdoors in windy, dry, cold, hot, air-conditioned, heated environments. Environment-related limitations affect ability to function and daily activities of work, social, physical (exercise), and general enjoyment of life.</p>	<p>54 Persistence of symptoms when mouth breathing Symptoms do not improve or resolve with mouth breathing.</p>	<p>55 General disability (<i>difficulty functioning, performing daily activities</i>) Symptoms and sequelae of ENS result in the patient being in a state of general disability.</p>	

*** A heart rate of 90 bpm being a norm for one patient can be a 50% increase in heart rate (tachycardia) for a patient with norm of 60 bpm.

PATHOPHYSIOLOGY

ENS is a complication of turbinate surgery, where the turbinates have been excised or damaged resulting in impairment or loss of nasal physiological functions. Total turbinate excision is the most frequent cause, but other procedures can cause it as well. All turbinate procedures have the potential to cause ENS if performed too aggressively. Removal or reduction of the inferior turbinates are more likely to result in ENS, than are the middle turbinates. Some turbinate procedures increase the risk of ENS, such as total turbinectomy or surface cautery²⁰.

Significant studies concerned with pathophysiology of ENS are those of EB Kern and SM Houser^{21,22}.

ENS is a multifactorial condition²³. Factors contributing to the ENS include: alteration of airflow pattern, reduced mucosal area, decreased humidification and increased warming, reduced nasal airflow resistance, loss of sensory, tactile and thermal receptors needed for inhaled air treatment, and neural damage^{24,25}. These changes are related to the alteration of pulmonary function²⁶.

It is not known how much turbinate tissue can be removed or damaged before causing ENS. Descriptors like “over-resection”, “aggressive”, or “excessive” populate the literature as a characterization of what level of excision or damage can result in ENS. However, these descriptors have not been quantified in any studies, nor are they described in any set of profession-

al guidelines. The majority of patients treated for ENS have experienced significant resections or reductions and/or had surgery involving mucosal-damaging techniques.

Cool thermoreceptors

Paradoxical obstruction is a central feature of ENS, where patients commonly suffer from nasal obstruction and a decreased nasal airflow sensation despite a patent airway. Since sensation of nasal patency is related to trigeminal activation of cool thermoreceptors through mucosal cooling, research focused on this issue has proposed a link to nasal cool thermoreceptors as a factor in ENS²⁷. Findings indicate that the perception of nasal airflow by means of mucosal cooling is a more significant factor in ENS than air temperature or nasal resistance are²⁸. As it applies to ENS, patients with wide nasal passages may produce less mucosal cooling²⁹.

For example, in the case of a total inferior turbinectomy (TIT), computational fluid dynamics (CFD) study has revealed that in addition to a decrease in nasal resistance, nasal heating and humidification efficiencies are reduced to 60%-80% of their pre-surgical measures (near 100%)³⁰. Additionally, the optimal nasal air-conditioning is necessary for pulmonary gas exchange in order to avoid desiccation and adhesion of the alveolar capillary bed³¹.

These receptors, along with other nasal receptors (mechanoreceptors, proprioceptors, and warm thermoreceptors), which cover the nasal epithelium, may be absent due to turbinate excision or damaged - either due to mucosal damage secondary to mucosal-damaging turbinate procedures or to mucosal atrophy.

Airflow

Computer airflow modeling studies show that there is significant airflow and aerodynamics disruption in the nose following resection of nasal turbinates³². It is thought that changes in aerodynamics play a central role in ENS. Turbulent airflow and decreased nasal resistance are characteristic of ENS. Turbinates act as resistors to, first, participate in the breathing function to expand the lungs, and, second, to direct the air in a streamlined, laminar pattern of airflow. Nasal resistance in the ENS nose demonstrates below normal values, and thus, the term “decreased nasal resistance” connotes below normal values of nasal resistance in both inspiration and expiration³³.

Both actions are integral to pulmonary function. Adequate nasal resistance is necessary to for sufficient fulfillment of respiratory function: optimizing alveolar ventilation, preventing alveolar collapse, and maintaining elasticity of the lungs. Too little nasal resistance (also mouth breathing) can result in micro areas of poor ventilation in the lungs³⁴.

In addition, decreased nasal resistance can also impair sensation of airflow. Excessive nasal emptiness and turbulence can increase drying of the nasal mucosa, having a downstream effect on nasal physiology.

In contrast to the normal nose, where airflow is evenly distributed, disruption of aerodynamics in the ENS nose leads to dyspnea and other undifferentiated breathing difficulties. Nasal resistance contributes approximately half the resistance of the airways. Even small changes in nasal resistance affect total respiratory function³⁵.

Dyspnea can also be explained by study results, which support the fact that mucosal cool thermoreceptors are connected to respiratory centers in the brain³⁶.

Dryness/mucosal atrophy

Progressive or non-progressive mucosal atrophy is quite common among ENS patients. Impaired mucosa regeneration due to mucosal atrophy or dysfunction leads to greater vulnerability of the nasal epithelium and impaired mucociliary transport. The moisture in the nose is reduced since the mucinous cells are impaired or missing, and nasal secretions decrease. Impaired local defenses against infections can cause mucosal swelling together with constant nasal inflammation and infections³⁷.

Mucociliary clearance

Diminished or absent mucociliary clearance and unusually thick mucus are common features of ENS. An impaired system can result in chronic sinusitis for some patients, particularly those with total turbinectomies. A study found that 50% of patients with total turbinectomies who had no sinusitis before developed chronic sinusitis following surgery³⁸.

Neuropathy

Neural damage is thought to be a contributing factor in ENS. Damaged nerve fibers from turbinate surgery may play a role, although there is little evidence specifically.

Additional risk factors for ENS have not been studied, but some factors have been proposed including: climatic conditions, an underlying connective tissue disorder, poor wound healing, poor regrowth of sensory nerves injured in turbinate surgery, aging mucosa, and others.

DIAGNOSIS

Diagnosis of ENS should be considered in any patient complaining of nasal dryness and breathing difficulties following turbinate surgery. Suspicion of ENS is not excluded in patients who have seemingly adequate turbinate volume or who do not appear to have nasal dryness upon endoscopic or nasal examination. Since ENS is multifactorial, multiple tests are necessary to confirm diagnosis of ENS.

ENS can be diagnosed both objectively and subjectively. Current diagnostic method relies on subjective reported symptomatology (SNOT-25) or ENS6Q and a positive “cotton test”. Radiographical and nasal endoscopy findings also contribute to the diagnosis. Nasal dryness shows a good correlation with the degree of turbinate tissue loss ³⁹.

Objective tests can include imaging, computational fluid dynamic (CFD) study, manometry, pathology (e.g., squamous metaplasia), and endoscopy. Many of these tests are easy to administer in the clinical setting.

While diagnosis has emphasized this current diagnostic method, other tools of diagnosis which are not routinely utilized should be considered. These tools are described below in this section.

Objective tests

CFD (aerodynamics)

CFD (Computational Fluid Dynamics) is the most advanced, precise, and objective test used in assessment of nasal aerodynamics. It computes the airflow pattern, unilateral airflow allocation, regional airflow distribution, nasal resistance, airflow velocity, aerodynamic load ^{****}, vorticity, and other parameters in different sections of the nasal cavity.

CFD studies show decreased (below normal) nasal resistance levels and diminished airflow rates with more chaotic streamlines in the inferior turbinate resected population ⁴⁰.

Rhinomanometry

While not definitive for ENS, rhinomanometry is useful to confirm nasal patency, and more importantly, confirm lower than normal rates of nasal resistance characteristic of ENS. While computational fluid dynamics (CFD) study may provide better data for ENS on nasal resistance, the availability of rhinomanometry as a clinical tool makes it a useful contributor to diagnosis.

^{****} Aerodynamic load - the distribution of pressure and shear stress represent a distributed load over the surface.

It should be noted, however, that rhinomanometry is not as useful to assess the patient's subjective nasal patency and may produce biased results for ENS patients.

Mean total resistance in normal subjects is .23 Pa/cm³/s, ranging between .15 Pa/cm³/s and .39 Pa/cm³/s⁴¹.

Computer Tomography

CT findings indicative of ENS are: mucosal thickening of the paranasal sinus lining and enlargement of the intranasal space with bowing of the nasal walls⁴². Two other radiographic findings may indicate ENS as a recent study found. According to a study, the mucosal thickness of both the central and posterior segments of the septum in ENS subjects is significant when compared to non-ENS subjects⁴³.

Cotton test

This is both a diagnostic test and a pre-therapeutic test. Cotton soaked in saline is placed in the nasal cavity in an area of deficit where an implant may be positioned. The cotton remains in place for about 30 minutes during which time the patient is asked to breathe comfortably. If there is subjective improvement in breathing, and a change in sensation or symptoms, this is considered a positive result and indicative of ENS. Patient report that they experience more relaxed breathing and a decrease in suffocation.

The cotton test may not be as useful for ENS-type patients who have seemingly adequate tissue, and may have suffered damaged turbinate mucosa. The utility of the cotton test remains to be validated⁴⁴.

Biopsy

Nasal biopsy may confirm evidence of squamous metaplasia of the nasal epithelium. Histological features are: loss of normal pseudostratified columnar epithelium, serous mucus and glandular atrophy, loss of cilia, loss of goblet cells, inflammatory cell infiltrate and endarteritis obliterans⁴⁵.

Biopsy can gauge mucosal condition, since mucosal metaplasia (and dryness) may not be visible upon exam. Metaplasia may be limited to patches of the epithelium (partial) or be more extensive depending on the state of the disease process. Squamous metaplasia may not be present in all ENS patients because some patients experience ENS symptoms weeks to months after the turbinate surgery, and metaplasia is suggestive of long-term dryness.

A positive finding in a patient complaining of ENS symptoms is strong diagnostic indicator of ENS. Further studies are needed to examine the correlation between biopsy findings and symptoms of ENS, and standardize pathology criteria⁴⁶.

Intranasal Schirmer test

Nasal dryness is one of the most reliable symptoms for ENS diagnosis.

The intranasal Schirmer test is used to assess nasal secretion and nasal mucosal moisture. Filter paper strips are placed bilaterally along the anterior septum for 10 minutes and the median wetting distance is evaluated. In a preliminary study to quantify nasal secretion, the median wetting distance of the normal (healthy, non-smokers) group was 10.3mm (range 3.6-35.0mm), and the smoking group was 8.4mm (range 2.5-28.0mm)⁴⁷.

The Schirmer test is already used widely in evaluation of dry eye to assess aqueous tear production and in evaluation of dry mouth to assess salivary gland hypofunction.

Endoscopy or nasal examination

Nasal endoscopy can be used for an assessment of the visual quality of the nasal mucosa. Mucosa may be dry and pale with crusting. The nasal epithelium may be fragile and bleed easily. Evidence of a turbinate reduction will be present and increased intranasal space may be present. Dryness can be confirmed on examination.

Saccharin Test

The saccharine test is used simple test of nasal mucociliary clearance (MCC). Mucociliary clearance sweeps the mucus layer toward the nasopharynx by ciliary action. Normal mucociliary clearance is less than 20 minutes. In the test, a saccharine particle is placed near the anterior portion of the inferior turbinate. If the patient takes longer to detect the sweetness than normal time, this is considered delayed or prolonged duration. This test is inexpensive and easy to perform⁴⁸.

Subjective Tests

Subjective tests include: SNOT-25, ENS6Q, and SNOT-55 - each as described in the **Symptoms** section above.

MANAGEMENT

At the present time, there is no cure for empty nose syndrome. However, in some cases symptoms can be managed or improved.

Conservative management

Conservative management does not differ substantially from that of secondary atrophic rhinitis and focuses on nasal hygiene and moisturizing nasal mucosa. Use of saline sprays and irrigations, gels, emollients, and humidifiers are meant to maintain the health of the remaining nasal mucosa and support the function of mucociliary clearance. Moisturizing nasal mucosa alleviates pain and discomfort from dryness, thick or absent secretions, prevents infection, reduces mucosal inflammation, irritation, improves breathing, as well as provides a protective barrier from the elements. Chronic dryness can lead to metaplasia, so it is important to keep the nose moist.

Patients need to be educated about the role of nasal hygiene and care and the use of topical agents in the management of ENS. Patient adjustment to the need for nasal care is often difficult – it is time consuming and requires a great deal of patient education. While conservative management should be encouraged, patients report that chronic use of saline spray and irrigations can be drying, and that gels and emollients can be irritating.

Moisture lasts minutes to a few hours, so patients are advised to repeat often as wished without generating too much additional irritation. ENS patients sometimes may have to use these tools sparingly so as not to worsen dryness or irritation. There is a role for topical corticosteroids and antibiotic ointment or therapy. Concern for lung health is important when selecting a product.

Frequency & duration of treatment

It may take many months of moisturizing to reach a level improvement where the patient is comfortable and no longer has severe burning, irritation, and pain. Moisturizing must continue regularly throughout the patient's lifetime. Patients with severe crusting may require removal of crusts by a physician. The short-lived effects of saline, irrigation, and emollients require the patient to repeat frequently in order to stay comfortable. Patients with thick mucus lodged deep in the nasal cavity that impedes breathing often need to remove mucus frequently throughout the day.

Basic aspects of care

Basic care may include:

- Daily nasal irrigations of saline and saline sprays;
- Emollient oils (e.g. sesame, rose geranium, etc.), ointments, and gels can provide some temporary relief in cases of severe dryness and crusts; oils containing menthol may increase sensation of nasal patency⁴⁹;
- Using air humidifier;
- Using a CPAP/APAP/AIRVO machine (preferably with an inbuilt humidifier);
- Using nasal filters or nose clips to create some resistance;
- Wearing protective masks (e.g. cold weather face mask or asthma mask) to protect respiratory system and lungs from the damaging effects of dry/cold air;
- Keeping in a humidified environment and avoiding windy, dry, cold, hot, air-conditioned, and heated environments (also environments with significant particulate smells, dust);
- Moving to countries/locations with warm and humid climate.

General Protocol for Nasal Moisturizing and Management of ENS

- Saline spray (preferably buffered)(e.g. Breathe-Ease® XL)
- Saline irrigation with or without steroid* (e.g. budesonide solution)
- Emollient oil (e.g. sesame or rose geranium oil), nasal ointment, cream, drops, or spray (e.g. Rinopanteina, Ponaris, Siccalix, Premarin, Nozoil, etc.)
- Antibiotic ointment (such as Mupirocin)

Select items with regard to irritation factor and effects on lungs.

*Patients report steroid solution is very drying. Long-term effects of corticosteroids in the ENS nose are unclear.

Advice on nasal hygiene and conservative management of ENS can be found in the book Having Nasal Surgery? Don't You Become an Empty Nose Victim!, written by ENS sufferer, Christopher Martin⁵⁰.

Saline Spray and Irrigations

Moisturizing the nose with saline spray or irrigations can help modestly. In the ENS nose, saline spray and irrigations provide only brief moisture, and while useful for nasal hygiene to clear out thick secretions, repeated use can wash away the natural nasal mucosal barrier, leading to further drying. Some patients find a pulsatile irrigator helpful rather than the neti-pot. Patients commonly find they must use saline sparingly so as to avoid further drying. It is important to clean out the irrigator regularly so as to avoid contamination of the machine, which can contribute to more sinus infections.

Irrigation solution

Normal saline rinse, can be isotonic or hypertonic. Some patients use Ringer's Lactate solution instead, as they find it is easier on the mucosa than regular saline. Some patients also prefer Breathe-Ease® XL nasal irrigation formula.

Spray

Normal saline spray can be isotonic or hypertonic. Some patients find hypertonic sprays helpful in dislodging thick mucus. Spray should preferably be buffered. As with nasal irrigation, patient may prefer Ringer's Lactate, or hyaluronic acid based nasal spray that they can purchase or make themselves. Some patients also prefer Breathe-Ease® XL.

Emollients, Gels and other preparations

Sesame oil based emollients have a long history in treating nasal dryness. Mayo Clinic offers a sesame and rose geranium oil preparation. Ponaris nasal emollient is also well known, but is recommended with reservation because of its iodine content and possible adverse effects in thyroid sensitive populations. Petroleum-based emollients should be avoided. Other popular preparations include: Rinopanteina, Siccalix, Nozoil, and Premarin cream. Preservative free

products are preferable. Preparations should be selected with regard to their pulmonary effects.

There are many preparations for nasal dryness on the market. Selection should be made with reference to lung health. Dexpanthenol and hyaluronic acid based preparations are newer to the market. Related studies on dexpanthenol indicate its efficacy in the dry nose.^{51 52}

Irritation and burning are common adverse effects of emollients, gels, and other preparations. Some patients find emollients further decrease nasal sensation and in general all preparations can make nasal breathing more difficult.

Humidifiers

Warm mist or cool mist humidifiers are helpful. Travel or personal humidifiers are available.

Antibiotics

Mupirocin antibiotic ointment is recommended. Other antibiotic treatment may be indicated for severe infection or chronic sinusitis. Antibiotic nasal irrigations are indicated in advanced forms of ENS where *Klebsiella ozena* or other colonizing bacteria may be present. For management of chronic sinusitis in ENS patients see “Chronic sinusitis” in the section **Complications**.

Steroids

Studies on intranasal corticosteroids indicate that their safety⁵³. However, their long-term effects on the mucosa of patients with ENS is not as clear. It is known that they can be damaging under certain circumstances. Temporary use of sprays and solutions added to saline irrigation for acute inflammation may be indicated.

Nasal airflow restriction

Nasal filters can be worn to restrict airflow and retain moisture. Sanispira brand nasal filters have been used by European ENS patients. Placing cotton in the nostrils or nasal cavity is not recommended because of possible aspiration.

Breathing

Menthol can improve nasal airflow sensation⁵⁴. Patients with significant dyspnea may see mild improvement with breathing retraining or speech therapy. More on breathing can be found in the section **Complications**, subsection **Anxiety and Depression**.

Sore throat

The throat may be dry and irritated as a result of surrounding dryness of nasopharynx and throat. Dryness and irritation can also be caused by nasal moisturizing preparations. Using a humidifier, remaining in a humid environment, and moisturizing nose with saline can improve dryness. Using a less irritating emollient and decreasing frequency of use of moisturizers may help. Lozenges (especially mentholated which can temporarily improve nasal airflow sensation) can provide temporary relief. Hot teas, gargling, and vaporizers are helpful.

Dry mouth

The mouth may be mildly dry as a result of surrounding dryness of nasopharynx and throat. Dry mouth may be exacerbated by oral breathing. Dental issues may develop secondary to dry mouth. Products on the market designed for dry mouth are helpful. Drinking fluids is also helpful.

Surgical Treatment (Implantation)

Surgical treatment for ENS using implants is appropriate for patients who have experienced significant turbinate resections. This treatment can be helpful in improving breathing, but there are limitations, especially with respect to function such as humidification, regeneration, or immune protection of the original mucosa. Since implants do not recreate the original nasal structure and aerodynamics, they may provide only limited improvements for a patient even if implanting can be helpful in some patients, especially those with substantial resections.

The surgical goal of implantation is to recreate the shape of the removed turbinate and thus regain some of the nose's capabilities to adequately pressurize, streamline, heat, humidify, filter, and sense the airflow. The idea is to re-establish nasal resistance, increase mucosal surface area, and restore nasal aerodynamics⁵⁵.

Surgical method involves narrowing the airway to provide more nasal resistance and directing the airflow to untouched tissue and in a more physiological way — either by increasing the size of partially resected turbinates with implant material or by creating what are called neo or pseudo turbinates. This is accomplished through submucosal implantation of graft material. The location of an implant should ideally re-create the natural airflow patterns within the nose⁵⁶.

For patients who are ENS-IT, the implantation is placed in the nasal septum or floor mucosa, or lateral wall. For ENS-MT, the implantation is placed in the septum. For ENS-type, it can be placed submucosally in the residual turbinate⁵⁷.

Current implantation materials include: acellular dermal matrix (ADM), autograft (auricular and costal cartilage), and hydroxyapatite⁵⁸. ADM may be used in its micronized form (e.g. Acell) and injected in the turbinate of patients with less severe volume deficit. Acell implants in combination with Adipose derived stem cells (ADSC) are being tried in certain centers.

Opinions differ on the length surgical treatment should be deferred - ranging from a few months to a year, in which time neural healing may take place^{59 60}.

While not all patients (properly assessed candidates) will derive benefit from surgical intervention for ENS, it does appear to result in clinical improvement and a positive association with quality of Life (QOL)⁶¹. Evaluations were focused on improvement on the SNOT questionnaire. A systematic review of surgical interventions indicated that the average total SNOT score improved after surgery, with an average improvement of 29.0 (standard deviation, 15.9) points. In addition, at least 50% of patients reported an improvement of 30 SNOT points or more⁶². The improvement needs to be clinically significant⁶³. Long-term studies on clinical outcome beyond a follow-up period of 48 months are needed⁶⁴.

Complications may include: donor site morbidity, infection, extrusion, and resorption⁶⁵.

Experimental Treatment (Regenerative Medicine)

Experimental treatment includes a number of regenerative medicine based treatments. These include intra-turbinal or intranasal injections of various combinations of platelet-rich plasma (PRP), platelet rich lipotransfer (PRL), extracellular matrix (ECM), adipose derived stem cells (ADSC), stromal vascular fraction (SVF), and adipose tissue (fat grafts). A few practitioners (ENT & plastic surgeons) offer these treatments in the United States, Germany, Spain, and Italy with results in improvement of mucosal function and turbinate morphology. Differentiated (expanded) ADSC treatment is undergoing trial in China. Similar developments are occurring in South Korea and India ^{66 67 68 69}. Duration of the positive effects of these treatments is unknown. For example, concerning fat grafts, the percentage of adipose tissue resorption and length of time the process takes in the turbinate has not been studied. Patients with larger resections may not benefit from fat grafts to the degree they might with conventional grafts if adequate turbinate volume cannot be created.

FURTHER ADVICE

Nasal valve collapse

Nasal valve collapse can be helped with Breathe-Right nasal strips and nasal cones. Tender noses from damaged nasal mucosa can make it difficult to use these products. Valve collapse can be evaluated by means of CT scan comparison and Cottle test, for example ⁷⁰.

Further surgery

Further surgery should be weighed very carefully. Surgery to repair nasal valve or other structural collapse may complicate ENS. For example, nasal valve repair may widen the airway increasing airflow and further drying the mucosa. Septoplasty may also result in a widened airway having greater drying effects. Surgically induced nerve damage may have a negative effect with respect to lack of airflow sensation. Repairs to septal perforations can be considered since they may contribute to nasal dryness. Sinus surgery, if necessary, may be considered, but should be approached conservatively so as to preserve remaining turbinate tissue and nasal structures. Further turbinate surgeries are not recommended.

Surgery on adjacent areas to the nose that may further open the airway (e.g. palate, tonsils) should be considered with regard to the possible negative effects on the ENS and breathing.

Environmental Limitations/Concerns

ENS patients experience an increase in symptoms in windy, dry, cold, hot, air-conditioned, or heated environments. For this reason, buildings with HVAC, and the outdoors pose potential difficulty for the ENS patient. In both indoor and outdoor environments, the addition of blowing air or windy conditions can quickly dry and irritate the mucosa. Particulate matter such as dust, household and food products in powder form, outdoor particulate such as yard refuse can penetrate unimpeded through the nasal cavity to the lungs causing choking and breathing difficulties. Thus, ENS patients may have environmental-related limitations. These limitations can affect the patient's daily activities, especially work, social, physical exercise, and general enjoyment of life. Since the nose can no longer respond normally to the environment, patients find they must locate or create an environment to suit their nose, and simply avoid difficult environments altogether, preferring to remain in their homes, where environmental conditions are able to be controlled.

Recommendations: Travel with portable humidifiers and a CPAP with a built-in humidifier. Apply emollients when needed to protect from the elements, heating and air conditioning. Cover the nose if all else fails.

Exercise

Aerobic (cardio) based exercise may be difficult because of patient breathing problems and may need to be approached in moderation or with light exercise. Strength training is a good alternative form of exercise. Swimming is popular among ENS patients.

Travel

Travelling may pose difficulties. Suggestions include portable humidifiers such as the personal warm mist humidifier MyPurMist (can be used with a car adaptor) and water bottle-based personal cool mist humidifiers (also can be used with a car adaptor).

COMPLICATIONS

Dry eyes

Dry eye and blepharitis can be treated by an ophthalmologist or dry eye specialist. Lid hygiene, compresses, and eye drops can lessen severity of dry eye and prevent recurrence of blepharitis. Keeping the nose moist can reduce the severity of dry eye.

Sleep

ENS patients experiencing sleep problems should be referred to a sleep specialist to have a sleep study performed. Many ENS patients have sleep-disordered breathing, including upper airway resistance syndrome (UARS), hypopnea, and apnea, including central sleep apnea. Sleep deprivation has a significant effect on the ability to learn, can have metabolic effects, impair memory. It can increase risk for development of type 2 diabetes and heart disease, and Alzheimer's disease. Using a humidifier when sleeping can assist with sleep. Some ENS patients find CPAPs with built-in humidifiers helpful. Medication may be necessary, but should be selected with regard to drying effects on the nasal and adjacent mucosa, which can worsen breathing and increase nasal and facial pain.

Pulmonary

ENS patients can be referred to pulmonology for treatment of asthma and other breathing-related concerns secondary to ENS ⁷¹. Patients presenting with breathing difficulties should routinely be referred for pulmonary evaluation. Evaluation which reveals an ENS-related complication can then be treated in conjunction with a pulmonologist. ENS pulmonary complications have not yet been specifically studied, but literature on unified airway disease may be instructive.

Patients with acute dyspnea may benefit from breathing retraining or speech therapy. Medication can be considered, but the drying effects of the selected medication must be assessed. Moist environments can lessen cold-air hyperventilation-provoked bronchoconstriction and coughing induced by the patient inspiring cold, dry air ⁷². Patients have reported that asthma medications are drying and can aggravate ENS symptoms.

Pain Management

Pain can include neuralgia and burning sensations, trigeminal nerve-related pain, and other facial and nasal pain. Patients with pain that is not alleviated or reduced through nasal moisturizing or by use of over-the-counter pain medications can be referred to pain medication specialists (algologists). How-

ever, alternative therapies should be tried before referral to pain management because of the risk of increasing nasal dryness and unwanted medication side-effects, especially if the patient is already being treated for other ENS complications with medication. Patients with pain related to the sphenopalatine ganglion may be improved with a nerve block ⁷³.

Chronic Sinusitis

A subset of ENS patients may suffer from sinusitis. Some patients develop sinusitis as a complication of ENS. Impaired defense mechanisms of the nose contribute to chronic sinusitis ⁷⁴.

Anxiety and Depression

Anxiety and depression are secondary symptoms that can occur as a result of ENS.

ENS patients commonly begin to experience symptoms of anxiety and depression secondary to the onset of ENS, having not had such symptoms prior to surgery. Anxiety can manifest as state of panic related to the patient's "fight-or-flight" response being triggered by breathing difficulties, primarily: paradoxical obstruction, lack of nasal airflow sensation, and disturbed nasal airflow pattern. Depression stems in large part from the fact that nasal function has been severely damaged and that there is no cure for ENS. Additionally, a decrease in quality of life, sleep deprivation, physical suffering, and a requirement for chronic nasal care play a substantial role. While there is some incidence of pre-existing anxiety and/or depression among a small subset of patients just as in any other condition, which can worsen symptoms, these pre-existing conditions do not play causal role in ENS. **Nasal breathing is an involuntary life-sustaining process, and, in the ENS patient, that process has been disrupted affecting physical and psychological regulation.**

Studies on surgical treatment for ENS indicate that there is improvement in anxiety and depression following successful surgical treatment. This suggests when the disease burden is reduced psychological symptoms improve or resolve ^{75 76}. However, not all patients are able to achieve improvement with surgical treatment. Clinical improvement following surgical treatment depends upon the type and extent of damage in each individual patient, as well as other factors.

There are no specific studies on the incidence of anxiety and depression caused by ENS. Studies on the incidence of anxiety and depression in chronic illness may be relevant.

It is important to recognize the grieving process in anxiety and depression caused by an irreversible surgical complication (ENS). Grief focuses on the loss of a key sensory organ and impairment of a life-sustaining respiratory function. Other contributors to anxiety and depression include: lack of recognition and support from friends, family, and medical professionals; lack of available treatments; absence of insurance coverage for current treatments; inability to work due to severity of symptoms, stress from lack of income (e.g. some ENS patients end up homeless); difficulty obtaining disability.

Psychological support should include specialists who are familiar with treating patients with chronic illness (also breathing disorders and pain). If therapy is insufficient, a combination of counseling and medication may be warranted for optimal benefit. If treatment with medications is indicated, care must be taken to select medications that do not increase dryness and worsen ENS symptoms, especially nasal burning, pain, dryness, and tightness. However, most classes of psychotropic medication are drying. Benzodiazepines are without this side effect, but carry the risks of dependence and addiction and so are not a long term solution.

Relaxation techniques such as progressive muscle relaxation can have modest benefit, but those focused on breathing will be difficult for the ENS patient due to their difficulties with nasal breathing. Breathing retraining and diaphragmatic breathing can be helpful in assisting patients who have difficulty regulating breathing. Mouth breathing is not a reasonable alternative, it does not resolve ENS.

Compromised quality of breathing has a direct effect on nervous system function and other body systems⁷⁷.

Patient care should be coordinated with ENT, primary physicians, and other specialists in the event of complications.

PROGNOSIS

ENS is a disabling condition having a severe impact on quality of life. Conservative, non-surgical, treatment methods can produce mild to moderate improvement in some symptoms, but have little effect on the main, breathing-related, symptoms. Surgical treatment with implants can improve symptoms to a greater degree, including breathing-related symptoms. However, normal nasal physiology is not restored, and absent functional and structural features of the nose cannot be recreated. The patient may be left with significant impairment despite the degree of improvement produced by implantation.

While implants can certainly improve the lives of patients, they are not a cure for ENS. A substantial recovery or cure can be achieved by means of regenerative medicine, when nasal structures are restored both functionally and structurally. Patients are in urgent need for a cure.

PREVENTION

New diagnostics like CFD and computer simulation are suggested. They can be used to identify patients who may benefit from turbinate surgery and assist in surgical planning⁷⁸. There is not really a reliable and objective means of selecting patients who will benefit⁷⁹.

Using conservative tissue sparing techniques, minimally invasive surgery with the goal to preserve mucosa are advised⁸⁰. Damage to the overlying mucosa should be avoided, particularly as applies to cauterization.

A number of studies have examined the various methods remarking on their safety and utility⁸¹. Submucosal reductions with a microdebrider, radiofrequency reduction or outfracture (lateralize) may reduce the risk of ENS by preserving the surface mucosa⁸². However, lack of evidence-level 2 studies on turbinate reduction, especially long-term studies, means that there may be no ideal technique that can be recommended⁸³. Histopathological studies are needed to supplement studies focused on clinical efficacy. For example, a study on medium to long-term histopathological changes after coblation of the inferior turbinate, the long-term histological effects of reduction were found to be similar to those induced by laser.

These included: significant fibrosis, glandular and venous sinusoid depletion, a significantly decreased intact pseudostratified, ciliated, columnar respiratory epithelium, and a significantly increased partial epithelial shedding⁸⁴.

Turbinate reductions should not create excess dilation in what in the normal nose is a continuous uniform slit space. Normal respiratory function requires a sufficiently narrow flow channel, where there is physiological nasal airflow and adequate contact between the air and the mucosal surface^{85 86}.

Contraindications include: aging mucosa which increases the risk of post-operative atrophic rhinitis⁸⁷.

Prior to surgery

Patient informed consent should include ENS as a risk associated with turbinate reduction, listing ENS as a rare, but possible complication.

REFERENCES

1. Houser SM (2014) Does the Method of Inferior Turbinate Surgery Affect the Development of Empty Nose Syndrome?. *J Otol Rhinol* 3:3. Characterizes ENS as a breathing disorder in which dryness may also be present.
2. Hol MKS, Huizing EH. Treatment of inferior turbinate pathology: a review and critical evaluation of the different techniques. *Rhinology* 2000;38:157–166.
3. Surface cautery by inducing mucosal damage to the epithelium, that may negatively affect nasal physiology. Passali's 2003 paper on the randomized clinical trial with a 6 year follow up on the treatment of inferior turbinate hypertrophy demonstrated that laser cautery, electrocautery, and cryotherapy produced mucosal damage that "strongly interferes with nasal physiology". Passali D, Passali FM, Damiani V, Passali GC, Bellussi L. Treatment of inferior turbinate hypertrophy: a randomized clinical trial. *Ann Otol Rhinol Laryngol* 2003; 112:683–8. See also: Houser SM (2014) Does the Method of Inferior Turbinate Surgery Affect the Development of Empty Nose Syndrome?. *J Otol Rhinol* 3:3. and Hol MKS, Huizing EH. Treatment of inferior turbinate pathology: a review and critical evaluation of the different techniques. *Rhinology* 2000;38:157–166.
4. Houser's term "multifactorial". Houser SM, Chaabra N, Empty Nose Syndrome In: Lalwani AK, Pfister MHF, eds. *Recent Advances in Otolaryngology Head & Neck Surgery, Volume 5*. 1st ed. Philadelphia: Jaypee Brothers Medical Publishers Ltd; 2016.
5. Barelli, P., Nasopulmonary Physiology. In: Timmons, BH, Ley, R. (Eds.), *Behavioral and Psychological Approaches to Breathing Disorders*. Plenum Press, New York, 1994.
6. Jiang C, Wang F, Chen K, Shi R. Assessment of Surgical Results in Patients With Empty Nose Syndrome Using the 25-Item Sino-Nasal Outcome Test Evaluation. *JAMA Otolaryngol Head Neck Surg*. 2014;140(5):453-458.
7. WHO Disability Assessment Schedule 2.0 (WHODAS 2.0).
8. Passali D, Lauriello M, Anselmi M, Bellussi L. Treatment of hypertrophy of the inferior turbinate: long-term results in 382 patients randomly assigned to therapy. *Ann Otol Rhinol Laryngol* June 1999 vol. 108 no. 6 569-575.
9. Houser SM, Chaabra N, Empty Nose Syndrome In: Lalwani AK, Pfister MHF, eds. *Recent Advances in Otolaryngology Head & Neck Surgery, Volume 5*. 1st ed. Philadelphia: Jaypee Brothers Medical Publishers Ltd; 2016.
10. Cullen KA, Hall MJ, Golosinskiy A. *Ambulatory Surgery in the United States, 2006*. National health statistics reports; no 11. Revised. Hyattsville, MD: National Center for Health Statistics. 2009. For the procedure: "Turbinectomy" (ICD-9-CM Code 21.9) numbering 196,000 in 2006. Also for the same year, 260,000 septoplasties, 257,000 sinus cases, 134,000 plastics cases, and 103,000 other cases.
11. Bhattacharyya N. Ambulatory sinus and nasal surgery in the United States: demographics and perioperative outcomes. *Laryngoscope*. 2010; 120:635–638.
12. Sozansky J, Houser SM. Pathophysiology of empty nose syndrome *Laryngoscope*. 2015 Jan;125(1):70-4.
13. Sozansky J, Houser SM. Pathophysiology of empty nose syndrome *Laryngoscope*. 2015 Jan;125(1):70-4.
14. Chhabra N, Houser S. The Diagnosis and Management of Empty Nose Syndrome. *Otolaryngol Clin N Am*. 42 (2009) 311 -330.
15. Chhabra N, Houser SM. The diagnosis and management of empty nose syndrome. *Otolaryngol Clin North Am*. 2009;42(2):311-330.
16. Velasquez N., Thamboo A, Habib AR, Huang Z, Nayak JV The Empty Nose Syndrome 6-Item Questionnaire (ENS6Q): a validated 6-item questionnaire as a diagnostic aid for empty nose syndrome patients. *Int Forum Allergy Rhinol* 2016 Aug 24.
17. Houser, as above, and other works. Kern: Moore EJ, Kern EB. Atrophic rhinitis: a review of 242 cases. *Am J Rhinol* 15: 355–361, 2001; Martin: Martin, C. *Having Nasal Surgery? Don't You Become an Empty Nose Victim!* Nashville, TN: Cold Tree Press; 2007.
18. Mangin D, Coste A, Zerah F, Béquignon E, Papon J, Devars du mayne M. Étude de la prévalence du syndrome d'hyperventilation chez les patients atteints d'un syndrome du nez vide. *Annales françaises d'Oto-rhino-laryngologie et de Pathologie Cervico-faciale*. 2014. Oct; 131 (4): A97.
19. Spodick DH. Normal sinus heart rate: appropriate rate thresholds for sinus tachycardia and bradycardia. *South Med J*. 1996 Jul;89(7):666-7. Spodick DH. Normal sinus heart rate:

- appropriate rate thresholds for sinus tachycardia and bradycardia. *South Med J*. 1996 Jul;89(7):666-7.
20. Houser SM, Chaabra N, Empty Nose Syndrome In: Lalwani AK, Pfister MHF, eds. *Recent Advances in Otolaryngology Head & Neck Surgery, Volume 5*. 1st ed. Philadelphia: Jaypee Brothers Medical Publishers Ltd; 2016
 21. Moore EJ, Kern EB. Atrophic rhinitis: a review of 242 cases. *Am J Rhinol* 15: 355–361, 2001.
 22. Sozansky J, Houser SM Pathophysiology of empty nose syndrome. *Laryngoscope*. 2015 Jan;125(1):70-4.
 23. Houser SM, Chaabra N, Empty Nose Syndrome In: Lalwani AK, Pfister MHF, eds. *Recent Advances in Otolaryngology Head & Neck Surgery, Volume 5*. 1st ed. Philadelphia: Jaypee Brothers Medical Publishers Ltd; 2016.
 24. Scheithauer MO. Surgery of the turbinates and “empty nose” syndrome. *GMS Curr Top Otorhinolaryngol Head Neck Surg*. 2010;9. And others.
 25. Houser SM. Surgical Treatment for Empty Nose Syndrome. *Arch Otolaryngol Head Neck Surg*. 2007;133(9):858-863.
 26. Coste A, Dessi P, Serrano E (2012) Empty nose syndrome. *Eur Ann Otorhinolaryngol Head Neck Dis* 129 (2): 93-97.
 27. Sozansky J, Houser SM. The physiological mechanism for sensing nasal airflow: a literature review. *Int Forum Allergy Rhinol*. 2014; 4: 834-8.
 28. Zhao K, Blacker K, Luo Y, Bryant B, Jiang J. Perceiving nasal patency through mucosal cooling rather than air temperature or nasal resistance. *PLoS One*. 2011;6(10):e24618.
 29. Zhao K, Jiang J, Blacker K, Lyman B, Dalton P, Cowart BJ, et al. Regional peak mucosal cooling predicts the perception of nasal patency. *Laryngoscope*. 2014;124(3):589–95.
 30. Dayal A, Rhee JS, Garcia GJM, Impact of Middle versus Inferior Total Turbinectomy on Nasal Aerodynamics Otolaryngology -- Head and Neck Surgery. September 2016; 155: 518-525. First published on May 10, 2016.
 31. Lindemann J, Keck T. Numerical simulation and nNasal air-conditioning. *GMS Curr. Top Otorhinolaryngol. Head Neck Surg*. 2010; 9: Doc08.
 32. See for example, Di, MY, Jiang, Z, Gao, ZQ, Li, Z, An, YR & Lv, W. 2013 Numerical simulation of airflow fields in two typical nasal structures of empty nose syndrome: A computational fluid dynamics study. *PLoS ONE* 8 (12).
 33. In the study, below normal values of nasal resistance were demonstrated in the numerical simulation airflow study of both a bilateral radical inferior turbinectomy (ENS-IT) and middle turbinectomy (ENS-MT). The below normal values were especially apparent in the ENS-IT model on both inspiration and expiration. For example, inspiration values: Nasal Resistance, 10-2 Pa·s/cm³ Normal = 2.19±1.09; ENS(MT) = 1.51±0.61; NES (IT) 0.84±0.29. Di, MY., Jiang, Z., Gao, ZQ, Li, Z, An, YR & Lv, W. 2013. Numerical simulation of airflow fields in two typical nasal structures of empty nose syndrome: A computational fluid dynamics study. *PLoS ONE* 8 (12).
 34. Barelli P., Nasopulmonary Physiology. In: Timmons, BH, Ley, R. (eds.), *Behavioral and Psychological Approaches to Breathing Disorders*. Plenum Press: New York; 1994.
 35. Corey JP, Sahin-Yilmaz, A. Assessment of Nasal Function. In: Snow JB, Wackym PA. eds. *Bal-lenger’s Otorhinolaryngology: Head and Neck Surgery* 17. Shelton, Connecticut: BC Decker Inc; 2009.
 36. Sozansky J, Houser SM. Pathophysiology of empty nose syndrome *Laryngoscope*. 2015 Jan;125(1):70-4. See esp. the comments on dyspnea in ENS.
 37. Scheithauer MO. Surgery of the turbinates and “empty nose” syndrome. *GMS Curr Top Otorhinolaryngol Head Neck Surg*. 2010;9.
 38. Berenholz L, Kessler A, Sarfati S, Eviatar E, Segal S (1998) Chronic sinusitis: a sequela of inferior turbinectomy. *Am J Rhinol* 12: 257-261.
 39. Hong HR, Jang YJ. Correlation between remnant inferior turbinate volume and symptom severity of empty nose syndrome. *Laryngoscope*. 2016 Jun;126(6):1290-5. (Epub 2015 Dec 21).
 40. Di, MY., Jiang, Z., Gao, ZQ, Li, Z, An, YR & Lv, W. 2013. Numerical simulation of airflow fields in two typical nasal structures of empty nose syndrome: A computational fluid dynamics study. *PLoS ONE* 8 (12).
 41. Corey JP, Sahin-Yilmaz, A. Assessment of Nasal Function. In: Snow JB, Wackym PA. eds. *Bal-lenger’s Otorhinolaryngology: Head and Neck Surgery* 17. Shelton, Connecticut: BC Decker Inc; 2009.
 42. Moore EJ, Kern EB. Atrophic rhinitis: a review of 242 cases. *Am J Rhinol* 15: 355–361, 2001.
 43. Thamboo A, Velasquez N, Ayoub N, Nayak JV. Distinguishing computed tomography findings in patients with empty nose syndrome. *Int Forum Allergy Rhinol*. 2016 Jul 13.
 44. S.C. Leong. The clinical efficacy of surgical interventions for empty nose syndrome: a systematic review. *Laryngoscope*, 125 (7) (2015), pp. 1557–1562.
 45. Moore EJ, Kern EB. Atrophic rhinitis: a review of 242 cases. *Am J Rhinol* 15: 355–361, 2001.
 46. Studies on histopathology in atrophic rhinitis may be applicable. See for example: Bist SS1, Bisht M, Purohit JP, Saxena R. Study of histopathological changes in primary atrophic rhinitis. *ISRN Otolaryngol*. 2011 Dec 26;2011:269479. See also, Singh I, Raizada RM, Chaturvedi VN, Jain SK, Ingole SN. Study of histopathological changes in atrophic rhinitis. *Indian J Otolaryngol Head Neck Surg*. 1999 Jan;51(1):21-4.
 47. Lindemann, J., Tsakiropoulou, E., Rettinger, G. et al. The intranasal Schirmer test: a preliminary study to quantify nasal secretion. *Eur Arch Otorhinolaryngol* (2014) 271: 2963.
 48. Deborah S, Prathibha KM (2014) Measurement of Nasal Mucociliary Clearance. *Clin Res Pulmonol* 2(2): 1019.
 49. Freund W, Wunderlich AP, Stocker T, Schmitz BL, Scheithauer MO. Empty nose syndrome: limbic system activation observed by functional magnetic resonance imaging. *Laryngo-scope*.2011;121(9):2019–25.

50. Martin, C. *Having Nasal Surgery? Don't You Become an Empty Nose Victim!* Nashville, TN: Cold Tree Press; 2007.
51. Hahn C, Böhm M, Allekotte S, Mösges R. Tolerability and effects on quality of life of liposomal nasal spray treatment compared to nasal ointment containing dexpanthenol or isotonic NaCl spray in patients with rhinitis sicca. *European Archives of Oto-Rhino-Laryngology*. 2013;270(9):2465-2472.
52. Sonnemann U, Scherner O, Werkhäuser N (2014) Treatment of rhinitis sicca anterior with ectoine containing nasal spray. *J Allergy (Cairo)* 2014:273219.
53. Pien, LC. How long can my patient use intranasal steroid sprays? December 2005 | Volume 72 | Number 12 | Pages 1079-1082. Department of Pulmonary, Allergy, and Critical Care Medicine. Retrieved from: <http://www.clevelandclinicmeded.com/medicalpubs/ccjm/december2005/pien.htm> Cleveland Clinic.
54. Freund W, Wunderlich AP, Stöcker T, Schmitz BL, Scheithauer MO. Empty nose syndrome: limbic system activation observed by functional magnetic resonance imaging. *Laryngoscope* 2011; 121:2019–2025.
55. Flint T. Computational investigation of empty nose syndrome, Center for Turbulence Research (Stanford), Annual Research Briefs, 2015.
56. Houser SM. Surgical Treatment for Empty Nose Syndrome. *Arch Otolaryngol Head Neck Surg*. 2007;133(9):858-863.
57. Houser SM. Surgical Treatment for Empty Nose Syndrome. *Arch Otolaryngol Head Neck Surg*. 2007;133(9):858-863.
58. Different materials have been used for nasal mucosal tissue expansion, including autologous materials (eg, bone, cartilage, muscle, and fat) and biomaterials (eg, Teflon [DuPont, Parkersburg, West Virginia], Plastipore [Xomed, Jacksonville, Florida], Bone Source [Orthofix, Huntersville, North Carolina], Gore-Tex [Newark, Delaware], Alloderm [Life Cell]) ; hydroxyapatite and b-Tricalcium Phosphate.
59. Kuan EC, Suh JD, Wang MB. Empty nose syndrome. *Curr Allergy Asthma Rep*. 2015 Jan;15(1):493.
60. Houser SM, Chaabra N, Empty Nose Syndrome In: Lalwani AK, Pfister MHF, eds. *Recent Advances in Otolaryngology Head & Neck Surgery, Volume 5*. 1st ed. Philadelphia: Jaypee Brothers Medical Publishers Ltd; 2016.
61. See for example: Houser SM. Surgical Treatment for Empty Nose Syndrome. *Arch Otolaryngol Head Neck Surg*. 2007;133(9):858-863. Also see, Jiang C, Wang F, Chen K, Shi R. Assessment of Surgical Results in Patients with Empty Nose Syndrome Using the 25-Item Sino-Nasal Outcome Test Evaluation. *JAMA Otolaryngol Head Neck Surg*. 2014;140(5):453-458, which evaluated Medpor implant surgery finding it had a positive association with QOL in patients with ENS, as seen in the differences in SNOT-25 scores before and after surgery. And also see, Tam YY, Lee TJ, Wu CC, Chang PH, Chen YW, Fu CH, Huang CC. Clinical analysis of submucosal Medpor implantation for empty nose syndrome. *Rhinology*. 2014 Mar;52(1):35-40, a study of patients who received Medpor implants saw a significant improvement of SNOT-22 pre-operatively to one year post-operatively.
62. S.C. Leong. The clinical efficacy of surgical interventions for empty nose syndrome: a systematic review. *Laryngoscope*, 125 (7) (2015), pp. 1557–1562.
63. Payne SC. Empty nose syndrome: What are we really talking about? *Otolaryngol Clin North Am* 42:331–337, ix-x, 2009.
64. S.C. Leong. The clinical efficacy of surgical interventions for empty nose syndrome: a systematic review. *Laryngoscope*, 125 (7) (2015), pp. 1557–1562.
65. Sharma A, Janus JR, Hamilton GS. *Regenerative Medicine and Nasal Surgery*. *Mayo Clin Proc*. 2015 Jan;90(1):148-58.
66. Friji, M.T., Gopalakrishnan, S., Verma, S.K., Parida, P.K. and Mohapatra, D.P. (2014), New regenerative approach to atrophic rhinitis using autologous lipoaspirate transfer and platelet-rich plasma in five patients: Our Experience. *Clin Otolaryngol*, 39: 289–292.
67. Balur MB, Oktay MF, Taskin U, Aydin S, Yucebas K, Altınay S, Celik DS. The Effect of Adipose Tissue on Atrophic Rhinitis: An Animal Model. *Otolaryngology -- Head and Neck Surgery* September 2014 151: P249.
68. Businco LDR, New technologies for the prevention and treatment of empty nose syndrome: minimally invasive and regenerative surgery with PRL. *Aesthetic Medicine* (2016) 2 (1) pp. 25-32. aestheticmedicinejournal.org [ISSN 2421-7115] In: *Aesthetic Medicine* 2 (1) pp. 25-32 aestheticmedicinejournal.org.
69. Xu X, Li L, Wang C, Liu Y, Chen C, Yan J, Ding H, Tang SY. The expansion of autologous adipose-derived stem cells in vitro for the functional reconstruction of nasal mucosal tissue. *Cell Biosci*. 2015 Sep 17;5:54.
70. A reformatting of CT scans can be performed to evaluate the nasal valve. See: Bloom JD, Sridharan S, Hagiwara M, Babb JS, White WM, Constantinides M. Reformatted computed tomography to assess the internal nasal valve and association with physical examination. *Arch Facial Plast Surg*. 2012 Sep-Oct;14(5):331-5. See also: Moche JA, Cohen JC, Pearlman SJ. Axial computed tomography evaluation of the internal nasal valve correlates with clinical valve narrowing and patient complaint. *Int Forum Allergy Rhinol*. 2013 Jul;3(7):592-7.
71. Works on Unified Airway Disease/Theory may be applicable to ENS, although there are no specific studies on the specific relationship. On unified airway disease, see for example, Meena RS, Meena D, Aseri Y, Singh BK, Verma PC. Chronic Rhino-Sinusitis and Asthma: Concept of Unified Airway Disease (UAD) and its Impact in Otolaryngology. *Indian Journal of Otolaryngology and Head & Neck Surgery*. 2013;65(Suppl 2):338-342.
72. Koskela HO (2007) Cold-air provoked respiratory symptoms: the mechanisms and management. *Int J Circumpolar Health* 66(2): 91–100.
73. Moore EJ, Kern EB. Atrophic rhinitis: a review of 242 cases. *Am J Rhinol* 15: 355–361, 2001.

74. Moore EJ, Kern EB. Atrophic rhinitis: a review of 242 cases. *Am J Rhinol* 15: 355–361, 2001.
75. Leong and others. S.C. Leong. The clinical efficacy of surgical interventions for empty nose syndrome: a systematic review. *Laryngoscope*, 125 (7) (2015), pp. 1557–1562.
76. Lee TJ, Fu CH, Wu CL, Tam YY, Huang CC, Chang PH, Chen YW, Wu MH. Evaluation of depression and anxiety in empty nose syndrome after surgical treatment. *Laryngoscope*. 2016 Jun;126(6):1284-9.
77. Courtney R (2009) The function of breathing and its dysfunctions and their relationship to breathing therapy. *International Journal of Osteopathic Medicine* 78–85.
78. Hariri BM, Rhee JS, Garcia GJ. Identifying patients who may benefit from inferior turbinate reduction using computer simulations. *Laryngoscope*. 2015 Dec;125(12):2635-41.
79. Roth M, and Kennedy DW. The case for inferior turbinate preservation. In: Tos M, Thomsen J, Balle V eds. *Rhinology: A State of the Art: Proceedings of the XVth Congress of the European Rhinologic Society and the XIIIth International Symposium on Infection and Allergy of the Nose: Copenhagen, Denmark, June 19-23, 1994*. New York: Kugler Publications; 1995.
80. Houser SM, Chaabra N, Empty Nose Syndrome In: Lalwani AK, Pfister MHF, eds. *Recent Advances in Otolaryngology Head & Neck Surgery, Volume 5*. 1st ed. Philadelphia: Jaypee Brothers Medical Publishers Ltd; 2016.
81. Hol MKS, Huizing EH. Treatment of inferior turbinate pathology: a review and critical evaluation of the different techniques. *Rhinology* 2000;38:157–166.
82. Houser SM, Chaabra N, Empty Nose Syndrome In: Lalwani AK, Pfister MHF, eds. *Recent Advances in Otolaryngology Head & Neck Surgery, Volume 5*. 1st ed. Philadelphia: Jaypee Brothers Medical Publishers Ltd; 2016.
83. Scheithauer MO. Surgery of the turbinates and “empty nose” syndrome. *GMS Curr Top Otorhinolaryngol Head Neck Surg*. 2010;9.
84. Berger G, Ophir D, Pitaro K, Landsberg R. Histopathological Changes After Coblation Inferior Turbinate Reduction. *Arch Otolaryngol Head Neck Surg*. 2008;134(8):819-823.
85. Mlynski G. Restorative procedures in disturbed function of the upper airways - nasal breathing. *GMS Current Topics in Otorhinolaryngology, Head and Neck Surgery*. 2005;4:Doc07.
86. Mlynski G. Physiology and Pathophysiology of Nasal Breathing. In: T. Metin Önerci ed, *Nasal Physiology and Pathophysiology of Nasal Disorders*. New York: Springer:2013.
87. Bofares KM. Dilemma of Inferior Turbinate Surgery. *Global Journal of Medical Research: Dentistry and Otolaryngology*. 2015;15(1) Version 1.0.t



EMPTY NOSE SYNDROME
INTERNATIONAL
ASSOCIATION